



Advancing the future of pediatric healthcare...together.

**Pediatric Otolaryngology** 

## Matthew t. Brigger, MD, MPH

1.	Activity Limitations:
	1. Keep the operated ear dry (especially no swimming)
	2. No nose blowing
	<ul> <li>Reduce Physical Activities</li> <li>PE restrictions: \( \frac{1}{2} \) Nonedays</li> <li>No school for days</li> </ul>
2.	Wound Care:  1. If present remove the ear dressing tomorrow.
	<ol><li>Cotton ball in ear- Change with a clean cotton ball 2-3 ting a day. Expect some bloody drainage on cotton ball several days. Stop using the cotton ball when there is more drainage coming out of the ear.</li></ol>
3.	Medications:
	1. Antibiotic
	2. Pain Medication
	<ul> <li>Tylenol or Ibuprofen is often all that is needed.</li> </ul>
	<ul> <li>A stronger pain medication may also be prescribed s as Norco or Hycet.</li> </ul>
	3. Ear Drops
	drops # of drops times a day d
	Begin
4.	<u>Diet</u> - Regular
5.	Follow-up visit

bleeding, severe pain, swelling, or redness behind the ear.

Division Chief Morgan Bliss, MD Daniela S. Carvalho, MD Wen Jiang, MD Shelby Leuin, MD Anthony E. Magit, MD, FAAP, MPH

Fellow:

Alex Manteghi, MD

Javan Nation, MD

Seth M. Pransky, MD, FAAP

Physician Assistant:

Denise M. Lago, PA-C

Main Office:

3030 Children's Way

Suite 112

San Diego, CA 92123

(858) 309-7701

(858) 966-8038 Fax

Regional Offices:

Encinitas

(760) 944-5545

Escondido

(760) 755-7600

Murrieta

(951) 600-1640

Oceanside

(760)547-1020

www.RCSSD.org

Revised 07-17-14