
Causes

You might get low blood sugar (also called hypoglycemia) if you:

- Take certain medicines and eat too few carbohydrates, or skip or delay a meal
- Take too much insulin or diabetes pills (ask your diabetes care team if this applies to you)
- Are more active than usual

Signs and Symptoms




If low blood sugar is not treated, it can become severe and cause you to pass out. If low blood sugar is a problem for you, talk to your doctor or diabetes care team.

Low blood sugar (Hypoglycemia)

What to do if you think you have low blood sugar

Check your blood sugar right away if you have any symptoms of low blood sugar. If you think your blood sugar is low but cannot check it at that time, treat anyway.



- 4 ounces (½ cup) of regular fruit juice (like orange, apple, or grape juice)
- 4 ounces (½ cup) of regular soda pop (not diet)
- 3 or 4 glucose tablets
- 5 to 6 hard candies that you can chew quickly (such as mints)

Wait 15 minutes and then check your blood sugar again. If it is still low, eat or drink something high in sugar again. Once your blood sugar returns to normal, eat a meal or snack. This can help keep low blood sugar from coming back.







For more information, visit Cornerstones4Care.com

Novo Nordisk Inc. grants permission to reproduce this piece for nonprofit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time.

Cornerstones4Care® is a registered trademark of Novo Nordisk A/S.

