Preparation Instructions for Colonoscopy With or without Upper Endoscopy (EGD)

Patient Weight: 68 lbs. to 76 lbs.

WHAT YOU WILL NEED FOR CLEAN OUT (All are over the counter medications)

MEDICATIONS:

MiraLAX OR Generic Polyethylene Glycol

Dulcolax (*tablets*) OR Generic Bisacodyl (*tablets*) OR Senna (*liquid*) CLEAR LIQUIDS: Water, tea, Gatorade, or sports drink to mix MiraLAX. AVOID anything RED, BLUE, or PURPLE - It could look like blood when it comes out as stool.



QUESTIONS OR PROBLEMS?

Call 858-966-4003 (option 3) If after 5 pm, call 858-576-1700 (option 0). Ask to page the on-call GI Fellow.

NEED TO CANCEL OR RESCHEDULE A PROCEDURE?

Please cancel or reschedule procedure(s) at least **72 hours before** your procedure date. Call 858-966-4003 (option 4) to speak with a scheduler

NO EATING SOLID FOOD ALL DAY.

CLEAR LIQUIDS ONLY (Liquids you can see through).

e.g., Ice popsicles, jello, soup broth, juice (no milk, orange juice, or shakes). <u>Gatorade or any</u> <u>sport drinks are preferred.</u>

AVOID anything **RED**, **BLUE**, or **PURPLE**.

CLEAN-OUT INSTRUCTIONS:

<u>At 8 am:</u>

Have your child take 3 tablet(s) of Dulcolax (Bisacodyl) OR 15 mL of Senna

1 hour after taking Dulcolax (Bisacodyl) OR Senna:

Mix **136** g (= **8** capfuls or packets) of MiraLAX (polyethylene glycol) with **64** ounces clear liquid. Have your child drink **6 to 7** ounces of this mixture every 15 to 20 minutes.

The Goal is to finish MiraLAX by 12:30 PM.

<u>After MiraLAX</u>, have your child drink 8 ounces of CLEAR LIQUIDS every hour. Preferably Gatorade or sports drink. The Goal is to drink 67 ounces (2 liters) before going to bed.

Your child can drink CLEAR LIQUIDS <u>until 2 hours before procedure</u>. STOP ALL LIQUIDS 2 HOURS BEFORE THE PROCEDURE. NO FOOD IS ALLOWED (NO EATING FOOD) ALL DAY and BEFORE THE PROCEDURE.

By 3 PM, if stools are NOT a clear liquid (See Great bowel prep photo below) Call 858-966-4003 (option 3) to discuss with GI nurse.

